



Low flow characteristics – the influence of geological conditions

Christian Reszler, Walter Poltnig, Till Harum, Peter Reichl

Institute of Water, Energy and Sustainability
Dept. of Water Resources Management
Joanneum Research, Graz, Austria


Objectives

- Analysis of geological influence on low flow characteristics (groundwater flow) in catchments
 - runoff components
 - water storage capacities
 - recession characteristics
 - subsurface drainage systemshydrographic vs. orographic catchment
- Case studies in catchments with comparable climatic conditions
 - Similarities and differences in runoff characteristics

Objectives


- Analysis of geological influence on low flow characteristics (groundwater flow) in catchments
 - runoff components
 - water storage capacities
 - recession characteristics
 - subsurface drainage systemshydrographic vs. orographic catchment
- Case studies in catchments with comparable climatic conditions
 - Similarities and differences in runoff characteristics
- Analysis assists in:
 - Evaluation of sensitivity of basins to dry climate conditions
 - Regionalisation
 - Modelling: structure and parameter identification

Case studies







Sattnitz







Sattnitz






Koralpe








Koralpe



Summit zone included in catchment (Schwanberg gauge)




Lower catchment (Hörmsdorf gauge)






Koralpe


Duration curves






Koralpe





Eastern Styria (Fluttendorf gauge)


Koralpe



Modelling

MIKE-SHE
(DHI)

conceptual approach
(system of linear
reservoirs)
→ parameters: time
constants and exchange
rates







Example Karawanken

Two neighbouring gauged catchments with differences in geology

Calibration






Example Karawanken


Two neighbouring gauged catchments with differences in geology

Calibration






Observed duration curves



Simulated runoff components (average 1985-2009)



Conclusions

Analysis of geological influence on low flow characteristics

- Evaluation of sensitivity to dry periods
 - long term water storage and GW recession:
weathered/fissured zones, karst, tectonics, geomorphology ...
(in combination with seasonality of recharge, e.g. snow melt,
autumn/winter rain storms → shift due to climate change)
 - large surface runoff intensifies vulnerability:
limited recharge, e.g. in tertiary regions

- Definition of hydrogeologically “homogenous” zones
for regionalisation of low flow characteristics
 - small scale heterogeneity in alpine region
 - data availability (monitoring network)

- Modelling: combined hydrological and geological analysis
assists in parameter identification and verification (plausibility
check) -> runoff components

Open research questions

- Local (small) scale → regional (large) scale:
 - regionalisation of low flow characteristics
 - hydrogeological unit subdivision: trade off
- Change of near surface weathering processes due to climate change:
 - soil
 - change of infiltration conditions
 - different vegetation
 - change in land use
 - ...

THANK YOU

Acknowledgements

- funded by: ETC PROJECT "ALP WATER SCARCE"
- Data and expertise:
Provincial governments of Carinthia and Styria




Photo: R. Benischke